

ON NILPOTENT GROUPS AND CONJUGACY CLASSES

EDITH ADAN-BANTE

ABSTRACT. Let G be a nilpotent group and $a \in G$. Let $a^G = \{g^{-1}ag \mid g \in G\}$ be the conjugacy class of a in G . Assume that a^G and b^G are conjugacy classes of G with the property that $|a^G| = |b^G| = p$, where p is an odd prime number. Set $a^G b^G = \{xy \mid x \in a^G, y \in b^G\}$. Then either $a^G b^G = (ab)^G$ or $a^G b^G$ is the union of at least $\frac{p+1}{2}$ distinct conjugacy classes. As an application of the previous result, given any nilpotent group G and any conjugacy class a^G of size p , we describe the square $a^G a^G$ of a^G in terms of conjugacy classes of G .

1. INTRODUCTION

Let G be a finite group, $a \in G$ and $a^G = \{a^g \mid g \in G\}$ be the conjugacy class of a in G . Let X be a G -invariant subset of G , i.e. $X^g = \{x^g \mid x \in X\} = X$ for all $g \in G$. Then X can be expressed as a union of n distinct conjugacy classes of G , for some integer $n > 0$. Set $\eta(X) = n$.

We can check that given any two conjugacy classes a^G and b^G of G , the product $a^G b^G = \{a^g b^h \mid g, h \in G\}$ is a G -invariant set. Thus $\eta(a^G b^G)$ is the number of distinct conjugacy classes of G such that $a^G b^G$ is the union of those classes. In this note we study the product $a^G b^G$ when G is a nilpotent group, and $|a^G| = |b^G| = p$ for some prime p . The main result is the following

Theorem A. *Let p be an odd prime number. Let G be a finite nilpotent group, a^G and b^G be conjugacy classes with the property that $|a^G| = |b^G| = p$. Then either $a^G b^G = (ab)^G$ or $a^G b^G$ is the union of at least $\frac{p+1}{2}$ distinct conjugacy classes.*

For a fix prime $p > 3$, we want to emphasize that Theorem A is a statement about a “gap” among the possible values that $\eta(a^G b^G)$ can take for any finite nilpotent group G and any conjugacy classes a^G and b^G of size p . We wonder if there exist any other “gaps”.

Given any prime $p > 2$ and any integer $n > 0$, we provide an example in Proposition 4.3 of a p -group G and a conjugacy class a^G of G such that $|a^G| = p^n$ and $\eta(a^G a^G) = \frac{p+1}{2}$. Thus the bound of Theorem A is optimal.

In Proposition 2.7 it is proved that if G is a finite group, a^G and b^G are conjugacy classes of size two, then $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) = 2$.

An application of Theorem A is the following

Theorem B. *Let p be an odd prime number. Let G be a nilpotent group and a^G be a conjugacy class of size p . Then one of the following holds*

i) $a^G a^G = (a^2)^G$. Therefore $[a, G] = [a^2, G]$ and $[a, G]$ is a normal subgroup of G .

Date: 2005.

1991 Mathematics Subject Classification. 20d15.

Key words and phrases. Finite p -groups, nilpotent groups, conjugacy classes.

ii) $a^G a^G$ is the union of exactly $\frac{p+1}{2}$ distinct conjugacy classes of G of size p .

Given any prime $p > 2$, we provide an example in Proposition 4.6 of a finite supersolvable group G and a conjugacy class a^G of G of size p with $\eta(a^G a^G) = 2$. Thus Theorem B may not remain true if G is only a supersolvable group.

For similar results in character theory, see [3].

Acknowledgment. I would like to thank Professor Everett C. Dade for very useful emails, and for helping me to improve the presentation of this note.

2. PROOF OF THEOREM A

Denote by $\mathbf{C}_G(a) = \{g \in G \mid a^g = a\}$ the centralizer of a in G . Also let $[a, g] = a^{-1}a^g$ and $[a, G] = \{[a, g] \mid g \in G\}$. Since $a^g = a[a, g]$ for any $g \in G$, we have that $a^G = \{ax \mid x \in [a, G]\}$. In particular, we get that $|a^G| = |[a, G]|$.

Through this note, we will use the well know fact that $|a^G| = |G : \mathbf{C}_G(a)|$.

Lemma 2.1. *Let p be an odd prime number, and r, s and t be integer numbers. The set $\{ri^2 + si + t \bmod p \mid i = 0, \dots, p-1\}$ has either one element or at least $\frac{p+1}{2}$ elements.*

Proof. If $r \equiv 0 \bmod p$, then $|\{ri^2 + si + t \bmod p \mid i = 0, \dots, p-1\}| = |\{si + t \bmod p \mid i = 0, \dots, p-1\}|$. If $s \equiv 0 \bmod p$, in addition, then $|\{si + t \bmod p \mid i = 0, \dots, p-1\}| = |\{t\}| = 1$. Otherwise $|\{si + t \bmod p \mid i = 0, \dots, p-1\}| = p$. Thus we may assume that $r \not\equiv 0 \bmod p$.

If $ri^2 + si + t \equiv rj^2 + sj + t \bmod p$, then $r(i^2 - j^2) \equiv r(i-j)(i+j) \equiv -s(i-j) \bmod p$. Therefore either $i - j \equiv 0 \bmod p$ or $r(i+j) \equiv -s \bmod p$. Thus either $i \equiv j \bmod p$ or $i \equiv -j - \frac{s}{r} \bmod p$. We conclude that the set $\{ri^2 + si + t \bmod p \mid i = 0, \dots, p-1\}$ either has only one element or it has at least $\frac{p}{2}$ elements, and therefore at least $\frac{p+1}{2}$ elements. $\square$

Lemma 2.2. *Let G be a finite p -group and N be a normal subgroup of G . Let a and b be elements of G . If $(aN)^{G/N} \cap (bN)^{G/N} = \emptyset$ then $a^G \cap b^G = \emptyset$. Thus $\eta((aN)^{G/N} (bN)^{G/N}) \leq \eta(a^G b^G)$.*

Proof. See Lemma 2.1 of [2]. $\square$

Lemma 2.3. *Let G be a finite group and $a, b \in G$. Then*

$$a^G b^G = ab[a^b, G][b, G].$$

Proof. See Lemma 2.1 of [1]. $\square$

Lemma 2.4. *Let G be a finite group, a^G and b^G be conjugacy classes of G . Assume that $|a^G| = |b^G| = |(ab)^G|$. Then $a^G b^G = (ab)^G$ if and only if $[ab, G] = [a, G] = [b, G]$ and $[ab, G]$ is a normal subgroup of G .*

Proof. Assume that $a^G b^G = (ab)^G$. By Lemma 2.3 we have that $[a^b, G][b, G] = [ab, G]$. Observe that $1_G \in [a^b, G]$ and $1_G \in [b, G]$. Observe also that $|[a^b, G]| = |[a, G]^b| = |[a, G]|$. Since $|a^G| = |b^G| = |(ab)^G|$, and $[a^b, G][b, G] = [ab, G]$, it follows that $[ab, G] = [a^b, G] = [b, G]$.

Let $u, v \in [ab, G]$. Since $[ab, G] = [a^b, G] = [b, G]$, we have that $u \in [a^b, G]$ and $v \in [b, G]$, and therefore $uv \in [a^b, G][b, G] = [ab, G]$. Since $1_G \in [ab, G]$ and G is a finite group, it follows that $[ab, G]$ is a subgroup of G . By Lemma 2.2 of [1] we have that $[ab, G]$ is a normal subgroup of G since it is a subgroup of

G . Thus $[a, G] = ([a, G]^b)^{b^{-1}} = [a^b, G]^{b^{-1}} = [ab, G]^{b^{-1}} = [ab, G]$ and therefore $[ab, G] = [a, G] = [b, G]$ is a normal subgroup of G .

If $[ab, G] = [a, G] = [b, G]$ and $[ab, G]$ is a normal subgroup of G , then $[a^b, G] = [a, G]^b = [a, G]$ and so $[a^b, G][b, G] = [ab, G]$. By Lemma 2.3 it follows that $a^G b^G = (ab)^G$ and the proof is complete. $\square$

Lemma 2.5. *Let G be a finite group, a^G and b^G be conjugacy classes of G . If $a^G b^G \cap \mathbf{Z}(G) \neq \emptyset$ then $\mathbf{C}_G(b) = \mathbf{C}_G(a^g)$ for some $g \in G$. If, in addition, $\mathbf{C}_G(a)$ is a normal subgroup of G then $\mathbf{C}_G(b) = \mathbf{C}_G(a)$.*

Proof. Let $z \in a^G b^G \cap \mathbf{Z}(G)$. Then $z = a^g b$ for some $g \in G$ and therefore $a^g = b^{-1}z$. Observe that

$$\begin{aligned} \mathbf{C}_G(a^g) &= \{h \in G \mid (a^g)^h = a^g\} = \{h \in G \mid (b^{-1}z)^h = b^{-1}z\} \\ &= \{h \in G \mid (b^{-1})^h z = b^{-1}z\} = \{h \in G \mid (b^{-1})^h = b^{-1}\} \\ &= \{h \in G \mid b^h = b\} = \mathbf{C}_G(b). \end{aligned}$$

We can check that $\mathbf{C}_G(a^g) = (\mathbf{C}_G(a))^g$ and the second statement follows. $\square$

Lemma 2.6. *Let G be a finite group, a^G and b^G be conjugacy classes of G with $\mathbf{C}_G(a) = \mathbf{C}_G(b)$ and $|a^G| = 2$. Then $\eta(a^G b^G) = 2$.*

Proof. See Proposition 3.4 of [1]. $\square$

Proposition 2.7. *Let G be a finite group, a^G and b^G be conjugacy classes of G with $|a^G| = |b^G| = 2$. Then either $a^G b^G = (ab)^G$ or $a^G b^G$ is the union of two distinct conjugacy classes of G .*

Proof. Since $|G : \mathbf{C}_G(a)| = |a^G| = 2$, we have that $\mathbf{C}_G(a) \triangleleft G$. If $a^G b^G \cap \mathbf{Z}(G) \neq \emptyset$ then by Lemma 2.5 we have that $\mathbf{C}_G(a) = \mathbf{C}_G(b)$. Thus by Lemma 2.6 we have that $\eta(a^G b^G) = 2$. We may assume then that $a^G b^G \cap \mathbf{Z}(G) = \emptyset$ and so if $c^G \subseteq a^G b^G$, then $|c^G| \geq 2$. Since $|a^G b^G| \leq 4$, it follows then that $\eta(a^G b^G) \leq 2$. $\square$

Lemma 2.8. *Let G be a finite group, b^G be a conjugacy class of G and $N \subseteq \mathbf{Z}(G)$. Then $b^G n$ is a conjugacy class of G for any $n \in N$. Also given any n_1 and n_2 in N , $b^G n_1 = b^G n_2$ if and only if $n_1 n_2^{-1} \in [b, G]$. Therefore, if $N \cap [b, G] = \{1_G\}$, then $b^G n_1 = b^G n_2$ if and only if $n_1 = n_2$.*

Proof. Since $N \subseteq \mathbf{Z}(G)$, $b^G n = (bn)^G$. Given any n_1, n_2 in N , observe that $b^G n_1 = b^G n_2$ if and only if $b^G = b^G n_1 n_2^{-1}$. Also $b^G = b^G n_1 n_2^{-1}$ if and only if there exist some $g \in G$ such that $b^g = b n_1 n_2^{-1}$, i.e. if and only if $n_1 n_2^{-1} \in [b, G]$. $\square$

Lemma 2.9. *Let K and L be finite groups, X be a K -invariant subset of K and Y be a L -invariant subset of L . Let $G = K \times L$ be the direct product of K and L . Set $X \times Y = \{(x, y) \in G \mid x \in X, y \in Y\}$. Then $X \times Y$ is a G -invariant subset of G and $\eta(X \times Y) = \eta(X)\eta(Y)$. In particular, given any conjugacy classes a^K and b^K of K , any conjugacy classes c^L and d^L of L , we have that $\eta((a, c)^G (b, d)^G) = \eta(a^K b^K) \eta(c^L d^L)$.*

Proof. Let $X = \cup_{i=1}^{\eta(X)} x_i^K$ and $Y = \cup_{j=1}^{\eta(Y)} y_j^L$. Since $G = K \times L$, given any $i \in \{1, \dots, \eta(X)\}$ and any $j \in \{1, \dots, \eta(Y)\}$, we can check that

$$(x_i, y_j)^G = \{(x, y) \mid x \in x_i^K, y \in y_j^L\} = (x_i^K, y_j^L).$$

Observe that $(x_i^K, y_j^L) = (x_{i_0}^K, y_{j_0}^L)$ if and only if $x_i^K = x_{i_0}^K$ and $y_j^L = y_{j_0}^L$. Since $X \times Y$ is the union of (x_i^K, y_j^L) for $i = 1, \dots, \eta(X)$, and $j = 1, \dots, \eta(Y)$, we conclude that $\eta(X \times Y) = \eta(X)\eta(Y)$. $\square$

Proof of Theorem A. Since p is a prime number and G is a nilpotent group, by Lemma 2.9 we may assume that G is a p -group.

Since $|a^G| = p$ and $|G : \mathbf{C}_G(a)| = |a^G|$, we have that $\mathbf{C}_G(a)$ is a normal subgroup of G of index p . Similarly, since $|b^G| = p$, we have that $\mathbf{C}_G(b)$ is a normal subgroup of G of index p . Also, since $\mathbf{C}_G(a) \cap \mathbf{C}_G(b) \leq \mathbf{C}_G(ab)$, it follows that $|G : \mathbf{C}_G(ab)| \leq p^2$.

Suppose that $|G : \mathbf{C}_G(ab)| = p^2$. Since $(ab)^G \subseteq a^G b^G$ and $|a^G| = |b^G| = p$, it follows then that $(ab)^G = a^G b^G$. Thus we may assume that $|G : \mathbf{C}_G(ab)| \leq p$ and so $|(ab)^G| \leq p$.

Let Z be a normal subgroup of G of order p contained in $\mathbf{C}_G(a)$. Observe such subgroup exists since $|\mathbf{C}_G(a)| > p$, otherwise G would have order p^2 and thus it would be abelian and therefore $|a^G| = 1$. Observe that Z is a central subgroup of G since G is nilpotent, Z is normal in G and $|Z| = p$. Fix $g \in G \setminus \mathbf{C}_G(a)$.

Claim 2.10. *If $|(aZ)^{G/Z}| = 1$ or $|(bZ)^{G/Z}| = 1$ then either $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) = p$.*

Proof. By symmetry, observe that it is enough to prove that if $|(aZ)^{G/Z}| = 1$ then either $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) = p$.

Observe that $|(aZ)^{G/Z}| = 1$ implies that $a^g = az$ for some $z \in Z \setminus \{1_G\}$, and therefore $[a, G] = Z$. Similarly, if $|(bZ)^{G/Z}| = 1$ then $[b, G] = Z$. Observe that then $|(aZ)^{G/Z}| = |(bZ)^{G/Z}| = 1$ implies that $a^G b^G = \{(ab)z \mid z \in Z\}$ and thus either $[ab, G] = Z$, and so $a^G b^G = (ab)^G$, or $[ab, G] \cap Z = \{1_G\}$. If $[ab, G] \cap Z = \{1_G\}$, by Lemma 2.8 we get that $\eta(a^G b^G) = |Z| = p$. We may assume then that $|(bZ)^{G/Z}| = p$ and thus $[b, G] \cap Z = \{1_G\}$. Since $a^g = az$, given any integer i we have that $a^{g^i} = az^i$. Thus

$$(2.11) \quad (ab)^G = \{ab[b, g^i]z^i \mid i = 0, \dots, p-1\},$$

and

$$(2.12) \quad a^G b^G = \{ab[b, g^i]z^j \mid i, j = 0, \dots, p-1\}.$$

Since $|(abZ)^{G/Z}|$ is either 1 or p , either $[ab, G] = Z$, and so $a^G b^G = (ab)^G$, or $[ab, G] \cap Z = \{1_G\}$. If $[ab, G] \cap Z = \{1_G\}$, by (2.11), (2.12) and Lemma 2.8 we get that $\eta(a^G b^G) = p$. Thus either $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) = p$. $\square$

By the previous claim, we may assume that $|(aZ)^{G/Z}| = |(bZ)^{G/Z}| = p$. Since $|a^G| = |b^G| = p$, we get that

$$[a, G] \cap Z = \{1_G\} \text{ and } [b, G] \cap Z = \{1_G\}.$$

By induction on the order of G , we have that either $\eta((aZ)^{G/Z}(bZ)^{G/Z}) = 1$ or $\eta((aZ)^{G/Z}(bZ)^{G/Z}) \geq \frac{p+1}{2}$. If $\eta((aZ)^{G/Z}(bZ)^{G/Z}) \geq \frac{p+1}{2}$, by Lemma 2.2 we have that $\eta(a^G b^G) \geq \frac{p+1}{2}$. We may assume then that $\eta((aZ)^{G/Z}(bZ)^{G/Z}) = 1$.

Observe that $|(abZ, G/Z)| > 1$ since $\eta((aZ)^{G/Z}(bZ)^{G/Z}) = 1$ and $|(aZ)^{G/Z}| = |(bZ)^{G/Z}| = p$. Thus $|(abZ, G/Z)| = p$. By Lemma 2.4 we have that $[aZ, G/Z] = [bZ, G/Z] = [(ab)Z, G/Z]$ is a normal subgroup of G/Z .

Let Y be a normal subgroup of G such that $Y/Z = [aZ, G/Z]$. Since $|[a, G]| = |(aZ)^{G/Z}| = p$, we have that Y is a normal subgroup of G of order p^2 and $[a, G]Z = Y$. Set $[a, g] = y$.

Claim 2.13. *If $\mathbf{C}_G(Y) \neq \mathbf{C}_G(a)$ or $\mathbf{C}_G(Y) \neq \mathbf{C}_G(b)$, then either $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) = p$.*

Proof. Since Y is normal in G , Z is central and $|Y : Z| = p$, either $|G : \mathbf{C}_G(Y)| = p$ or Y is contained in the center $\mathbf{Z}(G)$ of G .

Assume that $|G : \mathbf{C}_G(Y)| = p$ and $\mathbf{C}_G(Y) \neq \mathbf{C}_G(a)$. Fix $h \in \mathbf{C}_G(a) \setminus \mathbf{C}_G(Y)$. Observe such element exists since $\mathbf{C}_G(Y) \neq G$ and $\mathbf{C}_G(Y) \neq \mathbf{C}_G(a)$. Observe that $[h, y] \in Z \setminus \{1_G\}$. Set $[h, y] = z$. We can check that $[h^i, y] = z^i$ for $i = 0, \dots, p-1$. Since $[a, g] = y$, we have

$$(a^g)^{h^i} = (ay)^{h^i} = a^{h^i} y^{h^i} = ay z^i.$$

We can check also that $(a^{g^i})^{h^j}$ for $i, j = 1, 2, \dots, p$ are distinct elements. But then $|a^G| = p^2$. Thus, if $\mathbf{C}_G(a) \neq \mathbf{C}_G(Y)$ then $\mathbf{C}_G(Y) = G$.

Observe that $[b, g] \in Y$ since $[bZ, G/Z] = Y/Z$. Let $[b, g] = y^r z^s$ for some integers $r, s \in \{0, \dots, p-1\}$. We can check that then $[a, G] = \{y^i \mid i = 0, \dots, p-1\}$, $[b, G] = \{y^{ri} z^{si} \mid i = 0, \dots, p-1\}$, $[ab, G] = \{y^{(r+1)i} z^{si} \mid i = 0, \dots, p-1\}$ and $[a, G][b, G] = \{y^{ri+j} z^{si} \mid i, j = 0, \dots, p-1\}$. If $s \equiv 0 \pmod{p}$, then $[a, G][b, G] = [ab, G]$ and so $\eta(a^G b^G) = 1$. Otherwise by Lemma 2.3 we have that $a^G b^G = \cup_{v \in Z} (ab)^G v$, and since $[ab, G] \cap Z = \{1_G\}$, by Lemma 2.8 it follows that $\eta(a^G b^G) = p$.

Similarly, we can check that if $\mathbf{C}_G(Y) \neq \mathbf{C}_G(b)$, then either $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) = p$. $\square$

By the previous claim, we may assume that $\mathbf{C}_G(a) = \mathbf{C}_G(Y) = \mathbf{C}_G(b)$.

Set $[y, g] = z$. Since $g \in G \setminus \mathbf{C}_G(a)$, and by the previous claim it follows that $z \in Z \setminus \{1_G\}$. Because Z is central, $[y, g^i] = z^i$ for any integer i . Since $[a, g] = y$, we can check by induction on i that

$$(2.14) \quad a^{g^i} = ay^i z^{\frac{i(i-1)}{2}} \text{ for any integer } i \geq 0.$$

Since $a^{g^p} = a$ and $z^p = 1_G$, it follows that $y^p = 1_G$. Thus Y is an elementary abelian normal subgroup of G .

Let $b^g = by^r z^t$ for some integers r and t . Since $|b^G| = p$ and $[b, G] \cap Z = \{1_G\}$, we have that $r \not\equiv 0 \pmod{p}$. We can check by induction on i that

$$(2.15) \quad b^{g^i} = by^{ri} z^{r\frac{i(i-1)}{2} + ti} \text{ for any integer } i \geq 0.$$

Since $b \in \mathbf{C}_G(b) = \mathbf{C}_G(a) = \mathbf{C}_G(Y)$, combining (2.14) and (2.15), we get

$$(2.16) \quad (ab)^{g^i} = (ab)y^{(r+1)i} z^{(r+1)\frac{i(i-1)}{2} + ti}.$$

Since $[(ab)Z, G/Z] = [aZ, G/Z]$ and $|[aZ, G/Z]| = p$, we have that $[ab, G] \cap Z = \{1_G\}$. Thus we must have that $r+1 \not\equiv 0 \pmod{p}$. Fix $i \in \{0, \dots, p-1\}$. Let $j \in \{0, \dots, p-1\}$ such that $i \equiv (r+1)j \pmod{p}$. Then by (2.16) we have

$$(2.17) \quad ab^{g^i} = aby^i z^{\frac{i(i-1)}{2}} = (ab)^{g^j} z^{si},$$

where $s_i = \frac{i(i-1)}{2} - (r+1)\frac{j(j-1)}{2} - tj$. Set $S = \{s_i \mid i = 0, \dots, p-1\}$. Observe that

$$\begin{aligned} s_i &= \frac{i(i-1)}{2} - (r+1)\frac{j(j-1)}{2} - tj \\ &= \frac{(r+1)j((r+1)j-1)}{2} - (r+1)\frac{j(j-1)}{2} - tj \\ &= \frac{(r+1)r}{2}j^2 - tj. \end{aligned}$$

Since $r+1 \not\equiv 0 \pmod p$ and $i \equiv (r+1)j \pmod p$, $S \pmod p = \{s_i \pmod p \mid i = 0, \dots, p-1\} = \{\frac{r(r+1)}{2}j^2 - tj \pmod p \mid j = 0, \dots, p-1\}$. Therefore by Lemma 2.1 we have that either $S \pmod p$ has one element, or $S \pmod p$ has at least $\frac{p+1}{2}$. By (2.17), we have

$$(2.18) \quad a^G b^G = \cup_{i=0}^{p-1} (ab^{g^i})^G = \cup_{i=0}^{p-1} (ab)^G z^{s_i}.$$

Since $Z = \langle z \rangle$ is a central subgroup of G of order p , $[ab, G] \cap Z = \{1_G\}$, and the set $S \pmod p$ has either one element or at least $\frac{p+1}{2}$ elements, by Lemma 2.8 and (2.18) we have that either $\eta(a^G b^G) = 1$ or $\eta(a^G b^G) \geq \frac{p+1}{2}$ and the proof of the theorem is complete. $\square$

3. PROOF OF THEOREM B

The following two results are proved in [1].

Lemma 3.1. *Let G be a finite group, a^G and b^G be conjugacy classes of G . Assume that $\mathbf{C}_G(a) = \mathbf{C}_G(b)$. Then $a^G b^G = (ab)^G$ if and only if $[ab, G] = [a, G] = [b, G]$ and $[ab, G]$ is a normal subgroup of G .*

Lemma 3.2. *Let G be a group of odd order and a^G be the conjugacy class of a in G . Then*

$$(3.3) \quad \mathbf{Z}(G) \cap a^G a^G \neq \emptyset$$

if and only if $|a^G| = 1$. Thus if $|a^G| > 1$ and $b^G \subseteq a^G a^G$, then $|b^G| > 1$.

Proof of Theorem B. Since p is a prime number and G is a nilpotent group, by Lemma 2.9 we may assume that G is a p -group.

Since $|a^G| = p$ and $|a^G| = |G : \mathbf{C}_G(a)|$, $\mathbf{C}_G(a)$ is a normal subgroup of index p . Clearly $a \in \mathbf{C}_G(a)$, and since $\mathbf{C}_G(a)$ is normal in G , $a^g \in \mathbf{C}_G(a)$ for any $g \in G$. Thus $aa^g = a^g a$ for any $g \in G$.

Fix $g \in G \setminus \mathbf{C}_G(a)$. Since aa^{g^i} is G -conjugate to $a^{g^{p-i}} a$, $aa^{g^{p-i}} = a^{g^{p-i}} a$ and $a^G a^G = \cup_{i=0}^{p-1} (aa^{g^i})^G$, it follows that $\eta(a^G a^G) \leq \frac{p+1}{2}$. Thus by Theorem A we have that either $\eta(a^G a^G) = 1$ or $\eta(a^G a^G) = \frac{p+1}{2}$.

If $\eta(a^G a^G) = 1$, then by Lemma 3.1 we have that $[a, G] = [a^2, G]$ is a normal subgroup of G and i) holds.

We may assume now that $\eta(a^G a^G) = \frac{p+1}{2}$. Since G is of odd order, by Lemma 3.2, we have that $a^G a^G$ must be the union of conjugacy classes of size at least p . Since $|a^G a^G| \leq p^2$ and $\eta(a^G a^G) = \frac{p+1}{2}$, it follows that $a^G a^G$ is the union of $\frac{p+1}{2}$ distinct conjugacy classes of size p and ii) holds. $\square$

4. EXAMPLES

Hypotheses 4.1. Let p be an odd prime and $n > 0$ be an integer. Let G_0 be a p -group and $g_0^{G_0}$ be a conjugacy class of G with $g_0^{G_0} g_0^{G_0} = (g_0^2)^{G_0}$ and $|g_0^{G_0}| = p^{n-1}$.

Let $N = G_0 \times G_0 \times \cdots \times G_0$ be the direct product of p copies of G_0 . Set

$$a = (g_0, e, \dots, e),$$

where e is the identity of G_0 . Let $C = \langle c \rangle$ be a cyclic group of order p . Observe that C acts on N by

$$(4.2) \quad c : (n_0, n_1, \dots, n_{p-1}) \mapsto (n_{p-1}, n_0, \dots, n_{p-2})$$

for any $(n_0, n_1, \dots, n_{p-1}) \in N$.

Let G be the semi direct product of N and C , i.e G is the wreath product of G_0 and C .

Remark. Fix a prime $p > 2$ and an integer $l > 0$. Let E be an extraspecial group of exponent p and order p^{2l+1} . Let $g \in E \setminus \mathbf{Z}(E)$. We can check that $g^E = g\mathbf{Z}(E)$ and thus $g^E g^E = (g^2)^E$. Clearly if G is an abelian group and $g \in G$, then $g^G g^G = (g^2)^G$. Therefore given any prime p and any integer $l \geq 0$, there exist a p -group G and a conjugacy class a^G such that $a^G a^G = (a^2)^G$ and $|a^G| = p^l$, and so there always exists a group satisfying the previous hypothesis.

Proposition 4.3. Assume Hypotheses 4.1. Then $\eta(a^G a^G) = \frac{p+1}{2}$ and $|a^G| = p^n$. Thus given any integer $n > 0$ and any prime $p > 2$, there exist some p -subgroup G and some conjugacy class a^G of G such that $\eta(a^G a^G) = \frac{p+1}{2}$ and $|a^G| = p^n$.

Proof. Observe that $a^G = \{(x, e, \dots, e), (e, x, \dots, e), \dots, (e, \dots, e, x) \mid x \in g_0^{G_0}\}$. Thus $|a^G| = p|g_0^{G_0}| = p(p^{n-1}) = p^n$.

Observe also that

$$\begin{aligned} a^G a^G = & \{(xy, e, e, \dots, e)^b, (x, y, e, \dots, e)^b, (x, e, y, e, \dots, e)^b, \dots \\ & (x, e, \dots, e, y)^b \mid x, y \in g_0^{G_0} \text{ and } b \in C\}. \end{aligned}$$

Since $g_0^{G_0} g_0^{G_0} = (g_0^2)^{G_0}$, we have that

$$(4.4) \quad (a^2)^G = \{(x^2, e, \dots, e)^b \mid x \in g_0^{G_0} \text{ and } b \in C\}.$$

Set $a_i = (g_0, e, \dots, g_0, \dots, e)$, where the second g_0 is in the i -position, for example $a_0 = (g_0 g_0, e, \dots, e)$ and $a_1 = (g_0, g_0, e, \dots, e)$. Now observe that $(x, y, e, \dots, e)$ is G -conjugate to $(u, e, \dots, e, v)$ for any $x, y, u, v \in g_0^{G_0}$. Furthermore, we can check that $(x, e, \dots, e, y, e, \dots, e)$, where y is in the i -th position, is G -conjugate to $(u, e, \dots, v, e)$, where v is in the j -position if and only if $j \equiv \pm i \pmod{p}$. Thus $a_i^G \cap a_j^G \neq \emptyset$ if and only if $j \equiv \pm i \pmod{p}$. We can check that

$$a^G a^G = \cup_{i=0}^{\frac{p-1}{2}} a_i^G.$$

Therefore $\eta(a^G a^G) = 1 + \frac{p-1}{2} = \frac{p+1}{2}$. $\square$

Remark. Assume Hypotheses 4.1. Let $b = (g_0, g_0, e, \dots, e) \in N$. We can check that $\eta(a^G b^G) = p - 1$. The author wonders if there exist a prime $p > 5$, a p -group G , and conjugacy classes a^G and b^G of G with $|a^G| = |b^G| = p$ such that $\frac{p+1}{2} < \eta(a^G b^G) < p - 1$.

Hypotheses 4.5. Fix a prime p and let $F = \{0, 1, \dots, p-1\}$ be the finite field with p elements. Observe that F is also a vector space of dimension 1 over itself. Let $A = \text{Aff}(F)$ be the affine group of F . Observe that the group A is a cyclic by cyclic group and thus it is supersolvable.

Let C be a cyclic group of order p . Set $X = F$ and $K = C^X = \{f : X \rightarrow C\}$. Observe that K is a group via pointwise multiplication, and clearly A acts on this group (via its action on X).

Let G be the wreath product of C and A relative to X , i.e. $G = K \rtimes A$. We can check that G is a supersolvable group.

Proposition 4.6. Assume Hypotheses 4.5. Let e be the identity of C . Set $a = (c, e, e, \dots, e)$ in K , where $c \neq e$. Then $a \in G$, the conjugacy class a^G of G has size p and $a^G a^G = (a^2)^G \cup (c, c, e, \dots, e)^G$. Thus $\eta(a^G a^G) = 2$.

Therefore, given any prime p , there exist a supersolvable group G and a conjugacy class a^G of G with $|a^G| = p$ and $\eta(a^G) = 2$.

Proof. Observe that $a^G = \{(c, e, \dots, e), (e, c, e, \dots, e), \dots, (e, e, \dots, e, c)\}$. Thus a^G has p elements. We can check that

$$(4.7) \quad (a^2)^G = \{(c^2, e, \dots, e), (e, c^2, e, \dots, e), \dots, (e, e, \dots, e, c^2)\}.$$

Also, we can check that the set $(c, c, e, \dots, e)^G$ is the set of all elements of the form $(e, \dots, e, c, e, \dots, e, c, e, \dots, e)$, where the two c 's are in the i -th and the j -th position for some i, j with $1 \leq i < j \leq p$. Observe that

$$a^G a^G = \{(c^2, e, \dots, e), (e, c^2, e, \dots, e), \dots, (e, e, \dots, e, c^2)\} \cup (c, c, e, \dots, e)^G.$$

Thus $a^G a^G = (a^2)^G \cup (c, c, e, \dots, e)^G$. Clearly $(a^2)^G \neq (c, c, \dots, e)^G$. We conclude then that $\eta(a^G a^G) = 2$. $\square$

REFERENCES

- [1] E. Adan-Bante, Homogeneous products of conjugacy classes, to appear Arch. Math.
- [2] E. Adan-Bante, Conjugacy classes and finite p -groups, to appear Arch. Math.
- [3] E. Adan-Bante, Squares of characters and finite groups, preprint.

UNIVERSITY OF SOUTHERN MISSISSIPPI GULF COAST, 730 EAST BEACH BOULEVARD, LONG BEACH MS 39560

E-mail address: Edith.Bante@usm.edu